

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Note on Secchi's Companion to \$ 2481.

In 1856 Secchi found that the smaller star of the pair \$2481 was itself a close double. It was a very difficult object with his telescope, and he was able to measure it on only two nights. A few measures were made by other observers in the twenty-five years that followed, and these indicated, rather uncertainly, a slow retrograde motion. From 1881 on, the star seems to have been entirely neglected until 1897, when it was placed on my regular observing-list. I have followed it carefully for the past eight years, and can now say certainly that the fact of orbital motion is established and that the period is probably not much greater than fifty years.

Unfortunately the early measures are not only few in number but also very discordant. It is therefore impossible to compute even an approximate orbit at this time, though the observed arc is fully 340°. It will probably be necessary to wait at least twenty years for data to define the apastron end of the apparent ellipse.

The present note is written to call the attention of doublestar observers to this pair, which deserves annual measurement. So far as I can discover, the following list includes all the existing observations:—

MEASURES OF ₹ 2481 BC = SECCHI.

1856.832	93° ⋅4	0".4	$I^n$	Secchi.
59.610	98.2	0 .4	I	Secchi.
66.74	83 .4	o .45(es	st) I	O. STRUVE.
76.13	84 .8	0 .49	4	Schiaparelli.
<i>77</i> .31	69 .8	0 .37	2	Dembowski.
80.46	91 .1	0 .37	5	Schiaparelli.
81.57	61 .9	o .33	3	Hough.
97.85	243 .3	o .17	2	AITKEN.
98.60	2II .I	o .17	2	AITKEN.
1899.69	187 .9	0 .16	3	AITKEN.
1900.61	148 .2	0.15	I	AITKEN.
01.40	132 .8	0.14	2	AITKEN.
01.89	137 .1	0.15	1	AITKEN.
03.71	127 .2	0 .23	3	AITKEN.
04.44	120 .0	0 .22	2	AITKEN.
05.29	114 .3	0 .24	1	AITKEN.

May 22, 1905.

R. G. AITKEN.